

Postoperative Regime after Should

I a Subacromial Detrapment, Mobilization in Anesthesia

Free functional exercises

- Isometric muscle training in all directions
- Keep the thoraco-scapular sliding plane free, posture training
- Exercises to decrease joint capsular tension and stretching of the joint capsular
- Therapy of Myogelosis
- Passive/assistive movements in the range of motion

CPM if pain free

- Isotonic training increasing up to general muscular full load-bearing
- PNF
- Training with TheraBand, medical training therapy etc.
- Swimming exercises

I b LBT-Tenodesis / (Reversed-Endoprosthesis)

- 2 weeks shoulder sling, e. g. Medi Arm fix
- Passive/assistive movements in the range of motion
- Keep the thoraco-scapular sliding plane free, posture training
- Exercises to decrease joint capsular tension and stretching of the joint capsular

After 2 weeks:

Free functional exercises

- Isometric muscle training in all directions
- Keep the thoraco-scapular sliding plane free, posture training
- Exercises to decrease joint capsular tension and stretching of the joint capsular
- Therapy of Myogelosis
- Passive/assistive movements in the range of motion

CPM if pain free

- Isotonic training increasing up to general muscular full load-bearing
- PNF
- Training with TheraBand, medical training therapy etc.
- Swimming exercises

General Guidelines

- Active centering of the humerus head through the small shoulder muscles (pretension) before starting the specific exercise

Prof. Dr. Peter Strohm
Chief Physician

Prof. Dr. Volker Schöfl
Executive Physician
Shoulder Surgery & Sportorthopedics